

On the vapour trail of an atmospheric imprint in insects

M. D. Farnon Ellwood, Roger G. W. Northfield, Monica Mejia-Chang and Howard Griffiths

Biol. Lett. published online 16 February 2011
doi: 10.1098/rsbl.2010.1171

Supplementary data

["Data Supplement"](#)

<http://rsbl.royalsocietypublishing.org/content/suppl/2011/02/09/rsbl.2010.1171.DC1.html>

References

[This article cites 16 articles, 4 of which can be accessed free](#)

<http://rsbl.royalsocietypublishing.org/content/early/2011/02/09/rsbl.2010.1171.full.html#ref-list-1>

P<P

Published online 16 February 2011 in advance of the print journal.

Subject collections

Articles on similar topics can be found in the following collections

[ecology](#) (2156 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

On the vapour trail of an atmospheric imprint in insects

M. D. Farnon Ellwood^{1,3,*}, Roger G. W. Northfield¹,
Monica Mejia-Chang² and Howard Griffiths³

¹University Museum of Zoology and Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

²Global Ecology Unit, CREAF-CEAB-CSIC, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain

³Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, UK

*Author for correspondence (mdfe2@cam.ac.uk).

Terrestrial arthropods, at constant risk from desiccation, are highly sensitive to atmospheric temperature and humidity. A physiological marker of these abiotic conditions could highlight phenotypic adaptations, indicate niche partitioning, and predict responses to climate change for a group representing three-quarters of the Earth's animal species. We show that the ¹⁸O composition of insect haemolymph is such a measure, providing a dynamic and quantitatively predictable signal for respiratory gas exchange and inputs from atmospheric humidity. Using American cockroaches (*Periplaneta americana*) under defined experimental conditions, we show that insects respiring at low humidity demonstrate the expected enrichment in the ¹⁸O composition of haemolymph because of evaporation. At high humidity, however, diffusional influx of atmospheric water vapour into the animal forces haemolymph to become depleted in ¹⁸O. Additionally, using cockroaches sampled from natural habitats, we show that the haemolymph ¹⁸O signature is transferred to the organic material of the insect's exoskeleton. Insect cuticle, therefore, exhibits the mean atmospheric conditions surrounding the animals prior to moulting. This discovery will help to define the climatic tolerances of species and their habitat preferences, and offers a means of quantifying the balance between niche partitioning and 'neutral' processes in shaping complex tropical forest communities.

Keywords: stable isotopes; arthropods; niches; neutral theory; climate change

1. INTRODUCTION

In the face of global environmental change there is an urgent need to understand the rules governing the structure of ecological assemblages, and for unequivocal, quantitative evidence for the effects of climate disruption on the structure and dynamics of threatened communities [1]. At present, we lack a comprehensive understanding of the organization of ecosystems such as tropical forests, and of their gaseous exchanges with the atmosphere, which limits attempts to predict

the responses of these ecosystems to climate change [2]. Stable isotopes, widely used to answer intricate physiological and ecological questions in a variety of organisms [3], offer an elegant solution.

Insects and other tracheated arthropods, which make up three-quarters of all animal species [4], show a strong response to moisture gradients in tropical forests [5]. These animals respire through a series of spiracles, the tracheal openings which allow air to diffuse into the animal while optimizing gas exchange and controlling respiratory water loss [6]. As water evaporates, the lighter isotope (¹⁶O) changes phase and diffuses more rapidly than the heavier isotopomer (containing ¹⁸O). As a result, water vapour is depleted in ¹⁸O by around 10 parts per thousand (‰), whereas liquid water at the site of evaporation becomes enriched in ¹⁸O. The extent of ¹⁸O enrichment is controlled by the bidirectional exchange of water vapour between the surface of water bodies and the atmosphere, and is therefore a function of temperature and relative humidity (RH), as described by the Craig–Gordon equation [7].

Here, we demonstrate experimentally that the ¹⁸O signal of insect haemolymph in American cockroaches (*Periplaneta americana*) shows evaporative enrichment at low humidity, whereas under high humidity, there is an influx of ¹⁸O-depleted water vapour into the animal. Furthermore, the distinct signal correlating high humidity and water vapour uptake was present in the chitin of cockroaches from a range of natural habitats and contrasting global distributions.

2. MATERIAL AND METHODS

At the start of the experiment, 14 adult male cockroaches were placed into each of eight plastic chambers. Initial haemolymph ¹⁸O for these cockroaches was +2‰ versus Standard Mean Oceanic Water (SMOW) international isotopic standard, enriched compared with local tap water inputs (−8‰ versus SMOW) because of the low humidity (50% RH) in the controlled environment room. On days 2, 4 and 6, three cockroaches were removed from each chamber (the remaining five cockroaches were removed on day 8).

Water vapour was generated from two water sources, one 'enriched' in ¹⁸O ($\delta^{18}\text{O} = -8\text{‰}$: local, UK, tap water), the other relatively 'depleted' ($\delta^{18}\text{O} = -12\text{‰}$: Vienna, Austria, tap water). Two ambient RHs (98 and 50%) were derived by sparging air at 300 ml min⁻¹ through samples of each water source, maintained either at 30°C (saturated vapour pressure 3.71 kPa, to derive 98% RH) or 10°C (saturated vapour pressure 1.25 kPa, to derive 50% RH at the experimental chamber temperature). Half of the cockroaches (one set of four chambers) were provided with unlimited access to local, UK, tap water, $\delta^{18}\text{O} = -8\text{‰}$. Water consumption was measured daily using pipettes modified with a lateral orifice to allow insects to remove water ad libitum. In the second set of four chambers, cockroaches did not have access to drinking water.

Haemolymph ¹⁸O analysis was conducted indirectly by equilibrating a known concentration of CO₂ in the headspace of an Exetainer with 0.2 ml liquid for 3 days at 20°C, prior to purification and mass spectrometric analysis (VG Sira 10 Isotope ratio mass spectrometer). The precise ¹⁸O of Cambridge and Vienna water was $-8.41 \pm 0.26\text{‰}$ and $-12.49 \pm 0.21\text{‰}$ versus SMOW, and the ¹⁸O of water vapour generated from each source was $-18.23 \pm 0.27\text{‰}$ (Cambridge) and $-22.41 \pm 0.38\text{‰}$ (Vienna) versus SMOW. The equilibration and purification procedure were standardized on a daily basis against CO₂ equilibrated with a tap water laboratory standard and calibrated overall against two secondary standards [British Antarctic Survey (BAS) low and BAS high, respectively, -31.44 and -8.73‰ versus SMOW]. Chitin ¹⁸O analysis was completed on bulk samples from the hind tibia of five conspecific adult male cockroaches per habitat using a Thermo Finnigan 253 with TC (High Temperature Conversion) Elemental Analyser (Godwin Laboratory, University of Cambridge, UK) calibrated against NBS 127 and SMOW. These wild cockroaches were collected from two habitats per geographical location, preserved in alcohol and then dried.

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2010.1171> or via <http://rsbl.royalsocietypublishing.org>.

Received 11 December 2010
Accepted 21 January 2011






Figure 1. Effect of relative humidity (RH) and water vapour on the ^{18}O composition of the haemolymph of *P. americana*. (a) With drinking water at 50 versus 98% RH, while breathing enriched vapour at -18‰ from local UK tap water; (b) with drinking water at 50 versus 98% RH, while breathing depleted vapour at -22‰ from Vienna tap water; (c) without drinking water at 50 versus 98% RH, while breathing enriched vapour at -18‰ from local UK tap water; and (d) without drinking water at 50 versus 98% RH, while breathing depleted vapour at -22‰ from Vienna tap water (filled circles, observed; open circles, predicted).

To model the data, we used a parametrized non-steady state (NSS) version [8] of the Craig–Gordon enrichment equation (cf. equations 4 and 5 in Seibt *et al.* [8]). The NSS model integrates ^{18}O enrichment as a function of time and rate of (leaf) tissue-water turnover—or in this case—turnover of the cockroach haemolymph. Starting with the initial (control) $\delta^{18}\text{O}$ of insect haemolymph for each treatment, the model was parametrized with the respective $\delta^{18}\text{O}$ of water vapour, temperature, RH and measured weight change in each insect for the next 2 day sampling period. The NSS model was used to predict the haemolymph $\delta^{18}\text{O}$ for that subsequent period, and for each successive sampling period.

3. RESULTS

At low humidity (50% RH), both the measured data and the NSS model showed isotopic enrichment occurring in haemolymph through respiration and evaporative water loss (figure 1). However, at high atmospheric humidity (98% RH), water vapour uptake during respiration by *P. americana* reset the isotopic composition of the haemolymph, which became depleted ($\delta^{18}\text{O}$ more negative) as the influx of water vapour dominated the water budget of the insect (figure 1).

In cockroaches with access to drinking water, there was a delay in the isotopic depletion of the haemolymph compared with the model, but after six days

there was an agreement between modelled and observed values, with $\delta^{18}\text{O}$ decreasing to -9‰ in those cockroaches breathing enriched vapour (i.e. -1‰ below the ^{18}O composition of the drinking water; figure 1a), and -13‰ in the cockroaches breathing depleted vapour (i.e. -5‰ below the ^{18}O composition of the drinking water; figure 1b).

In cockroaches without access to drinking water, there was close agreement between observed and modelled values: after eight days $\delta^{18}\text{O}$ decreased to -11‰ in the cockroaches under enriched vapour, and to -16‰ in the cockroaches breathing depleted vapour (figure 1c,d). Interestingly, it seems that water vapour uptake and condensation can occur beyond normal equilibrium exchange, as seen by the decrease in haemolymph $\delta^{18}\text{O}$ below the value predicted by the model (figure 1d).

We compared the chitin from *P. americana* used experimentally with cockroach species collected from a variety of natural habitats, including the internal chambers of an epiphytic bird's nest fern (*Asplenium nidus*) in lowland tropical forest in Borneo [9]. There were significant differences in the ^{18}O composition of all species (one-way ANOVA: $F_{6,28} = 35.51$, $n = 35$,




Figure 2. Effect of RH on the ^{18}O composition of the chitin of cockroaches from natural habitats. Solid points show, in ascending order of latitude, the ^{18}O composition and the within-habitat RH for: (1) *Namablatta bitaeniata* and (2) *Tivia* sp. from the Brandberg, Swakopmund District, Namibia, $21^{\circ}9' \text{ S}$, $14^{\circ}35' \text{ E}$; (3) *Periplaneta americana*, our model insects, which were grown in controlled environment chambers in Cambridge, UK; (4) *Ischnoptera* sp. and (5) *Epilampra* sp. from the Refugio Los Volcanes, Santa Cruz, Bolivia, $18^{\circ}06' \text{ S}$, $63^{\circ}36' \text{ W}$; (6) *Margatta* sp. and (7) *Haania* sp. from inside an epiphytic bird's nest fern in Danum Valley, Sabah, Borneo, $4^{\circ}58' \text{ N}$, $117^{\circ}48' \text{ E}$ (mean \pm s.e., $n = 5$ cockroaches). Open points show the ^{18}O composition of the precipitation, and the mean local RH, at each location (mean \pm s.e., $n = 5$ recordings).

$p < 0.001$; figure 2). Most striking was the highly significant correlation between the depletion of ^{18}O in insect chitin and increasing RH (Pearson's $r = -0.989$, $p < 0.001$; solid symbols, figure 2). This relationship is apparent for both RH measured at the point of collection of each species (solid symbols, figure 2), and mean local RH calculated for the date of collection from the local air temperature and vapour pressure [10] (open symbols, figure 2). By contrast, the ^{18}O signatures of local precipitation likely to be ingested by the cockroaches [10] showed no significant relationship with the ^{18}O signature of cockroach chitin (Pearson's $r = -0.137$, $p = 0.769$; figure 2).

4. DISCUSSION

Insect haemolymph represents an isolated pool of water, and hence the ^{18}O isotopic composition can be defined by the Craig–Gordon model [7], which incorporates kinetic and equilibrium isotope transformations driven by temperature and humidity. Although the NSS version used in this paper was developed for use in plants [11], we hypothesized that short-term disequilibria in daily plant-water supplies and stomatal apertures might be equivalent to intermittent drinking and discontinuous gas exchange in insects [6], thus allowing the extent of water vapour uptake at high humidity to be tracked (cf. Helliker & Griffiths [11]). Source precipitation or ground-water determines the starting point for plant tissue-water and insect haemolymph signals, refined by temperature and humidity (see progression in figure 1). The integration of daily mean humidity is then transferred isotopically into the chitin ($\text{C}_8\text{H}_{13}\text{O}_5\text{N}$), together with a systematic (and constant) enrichment factor when the insect moults [12],

as we have shown for individual cockroaches from natural habitats, including those specifically associated with epiphytic bird's nest ferns in a rainforest canopy in Borneo (figure 2). Arthropods of this particular forest show a strong response to temperature and moisture gradients [5], which are a common feature of tropical forest canopies, often being associated with vertical stratification [2]. However, bird's nest ferns buffer microclimatic fluctuations [13], and provide discrete habitats for their invertebrate inhabitants [9]. We therefore propose that the ^{18}O signals in terrestrial arthropods can be used to integrate species distributions with microclimate and climate change [14], and that natural microcosms such as epiphytes offer the prospect for determining the extent to which niche partitioning, as compared with 'neutral' processes, shape complex tropical forest communities [15].

The experimental data show the interplay between source-water inputs and atmospheric humidity for insects: under low humidity, evaporative enrichment occurs, but under high humidity the ingested water $\delta^{18}\text{O}$ signal is reset by the isotopically depleted vapour influx (figure 1*a,b*), despite net evaporation of haemolymph (see the electronic supplementary material results and figure S1). Accordingly, the $\delta^{18}\text{O}$ of haemolymph was lower than that of drinking water by up to 5‰, particularly in those cockroaches breathing depleted vapour (-8‰ versus -13‰ ; figure 1*b*). This helps to explain the lack of correlation between the ^{18}O signals in cockroach chitin and local precipitation in South America, Africa and Borneo (figure 2), also seen in other studies [12].

With regard to the depletion of haemolymph $\delta^{18}\text{O}$ below that predicted by the model (figure 1*d*), it is possible that condensation is occurring inside the tracheoles, where water vapour might not be at isotopic

exchange equilibrium with the external atmosphere. Thus, when spiracles are closed, a Rayleigh distillation [16] could lead to the progressive condensation of lighter isotopes—similar to that seen in clouds with altitude—which would lead to progressive depletion of residual water vapour within the tracheoles, and haemolymph signals that are more depleted than predicted by the model. Meanwhile, whether there are isotopic shifts associated with water vapour absorption [17], or with fluid reabsorption by the Malpighian tubules [18], and whether the haemolymph of insects with contrasting life cycles responds in the same way as in cockroaches, has yet to be resolved.

The Departments of Zoology and Plant Sciences, University of Cambridge, supported this work. Darren Mann of the Oxford University Museum of Natural History kindly donated cockroach specimens. M.D.F.E. is grateful to Daphne Ellwood for her support and to Michael Akam, Adrian Friday and Simon Maddrell for their valuable discussion.

- 1 Pimm, S. L. 2009 Climate disruption and biodiversity. *Curr. Biol.* **19**, 595–601. (doi:10.1016/j.cub.2009.05.055)
- 2 Ozanne, C. M. P. *et al.* 2003 Biodiversity meets the atmosphere: a global view of forest canopies. *Science* **301**, 183–186. (doi:10.1126/science.1084507)
- 3 West, J. B., Bowen, G. J., Cerling, T. E. & Ehleringer, J. R. 2006 Stable isotopes as one of nature's ecological recorders. *Trends Ecol. Evol.* **21**, 408–414. (doi:10.1016/j.tree.2006.04.002)
- 4 Stork, N. E. 1988 Insect diversity: facts, fiction and speculation. *Biol. J. Linn. Soc.* **35**, 321–337. (doi:10.1111/j.1095-8312.1988.tb00474.x)
- 5 Dial, R. J., Ellwood, M. D. F., Turner, E. C. & Foster, W. A. 2006 Arthropod abundance, canopy structure, and microclimate in a Bornean lowland tropical rain forest. *Biotropica* **38**, 643–652. (doi:10.1111/j.1744-7429.2006.00181.x)
- 6 Schimpf, N. G., Matthews, P. G. D., Wilson, R. S. & White, C. R. 2009 Cockroaches breathe discontinuously to reduce respiratory water loss. *J. Exp. Biol.* **212**, 2773–2780. (doi:10.1242/jeb.031310)
- 7 Craig, H. & Gordon, L. 1965 Deuterium and oxygen 18 variations in the ocean and marine atmosphere. In *Stable isotopes in oceanographic studies and paleotemperatures* (ed. E. Tongiorgi), pp. 9–130. Pisa, Italy: Consiglio Nazionale Delle Ricerche Laboratorio di Geologia Nucleare.
- 8 Seibt, U., Wingate, L., Berry, J. A. & Lloyd, J. 2006 Non-steady state effects in diurnal ^{18}O discrimination by *Picea sitchensis* branches in the field. *Plant Cell Environ.* **29**, 928–939. (doi:10.1111/j.1365-3040.2005.01474.x)
- 9 Ellwood, M. D. F. & Foster, W. A. 2004 Doubling the estimate of invertebrate biomass in a rainforest canopy. *Nature* **429**, 549–551. (doi:10.1038/nature02560)
- 10 IAEA/WMO 2006 Global network of isotopes in precipitation. The GNIP database. See <http://www.iaea.org/water>.
- 11 Helliker, B. R. & Griffiths, H. 2007 Toward a plant-based proxy for the isotope ratio of atmospheric water vapour. *Global Change Biol.* **13**, 723–733. (doi:10.1111/j.1365-2486.2007.01325.x)
- 12 Schimmelmann, A. & DeNiro, M. J. 1986 Stable isotopic studies on chitin. III. The D/H and $^{18}\text{O}/^{16}\text{O}$ ratios in arthropod chitin. *Geochim. Cosmochim. Acta* **50**, 1485–1496. (doi:10.1016/0016-7037(86)90322-4)
- 13 Turner, E. C. & Foster, W. A. 2006 Assessing the influence of bird's nest ferns (*Asplenium* spp.) on the local microclimate across a range of habitat disturbances in Sabah, Malaysia. *Selbyana* **27**, 195–200.
- 14 McGill, B. J. 2010 Matters of scale. *Science* **328**, 575–576. (doi:10.1126/science.1188528)
- 15 Ellwood, M. D. F., Manica, A. & Foster, W. A. 2009 Stochastic and deterministic processes jointly structure tropical arthropod communities. *Ecol. Lett.* **12**, 277–284. (doi:10.1111/j.1461-0248.2009.01284.x)
- 16 Scott, K. M., Lu, X., Cavanaugh, C. M. & Liu, J. S. 2004 Optimal methods for estimating kinetic isotope effects from different forms of the Rayleigh distillation equation. *Geochim. Cosmochim. Acta* **68**, 433–442. (doi:10.1016/S0016-7037(03)00459-9)
- 17 O'Donnell, M. J. 1977 Site of water vapor absorption in the desert cockroach, *Arenivaga investigata*. *Proc. Natl Acad. Sci. USA* **74**, 1757–1760. (doi:10.1073/pnas.74.4.1757)
- 18 O'Donnell, M. J. & Maddrell, S. H. P. 1995 Fluid reabsorption and ion transport by the lower Malpighian tubules of adult female *Drosophila*. *J. Exp. Biol.* **198**, 1647–1653.